

**Appendix Table 1. Factors Independently Associated with Recurrent Pneumonia:
Multivariable Logistic Regression Analysis**

	Adjusted Odds Ratio (95% CI)	P- Value
Age >75 years	1.19 (0.90-1.57)	0.23
Male Sex	0.94 (0.72-1.22)	0.62
Age Subtracted Pneumonia Severity Index	1.00 (0.99-1.01)	0.99
<i>(per point)</i>		
Inhaled Corticosteroids	1.31 (0.93-1.83)	0.11
Not Fully Independent	1.61 (1.21-2.14)	0.001
<i>(Any Impairment in Functional Status)</i>		

**Appendix Table 2. Factors Independently Associated with Recurrent Pneumonia:
Multivariable Competing Risks Proportional Hazards Analysis**

	Adjusted Competing Risks Hazards Ratio (95% CI)	P- Value
Age >75 years	1.17 (0.89-1.53)	0.26
Male Sex	0.95 (0.73-1.22)	0.68
Age Subtracted Pneumonia Severity Index (<i>per point</i>)	1.00 (0.99-1.01)	0.99
Inhaled Corticosteroids	1.30 (0.95-1.79)	0.10
Not Fully Independent (<i>Any Impairment in Functional Status</i>)	1.58 (1.20-2.08)	0.001